

# Anti-Mouse IgM(μ chain specific), AlpSdAbs® VHH(HRP)

## Summary

Code	001-107-005
Immunogen	Recombinant Fc region of mouse IgM
Host	Alpaca pacous
Isotype	VHH domain of alpaca IgG2b/2c
Conjugate	HRP
Specificity	Fc region of mouse IgM(μ chain specific)
Cross-Reactivity	No cross-reactivity with mouse, rabbit, human, cynomolgus, rat, goat IgG Affinity: <1nM
Purity	Recombinant Expression and Affinity purified
Size	100ug
Buffer	10mM PBS (pH 7.5), 0.05% sucrose, 0.1% trehalose, 0.01% proclin300
Formation	Freeze-dried Powder, Store at 2-8 °C
Rehydration	Rehydrated by 100uL dH2O or 100uL dH2O with 50% Glycerol (Final concentration:1mg/mL)
Storage after Rehydration	-20 °C(Avoid freeze/thaw cycles), Stable for 12 months at -20°C, Protect from light

## Description

Anti-Mouse IgM(μ chain specific), AlpSdAbs® VHH(HRP) is designed for detecting mouse IgM specifically. Anti-Mouse IgM(μ chain specific), AlpSdAbs® VHH(HRP) is based on monovalent, recombinant single domain antibodies to mouse IgM(μ chain specific) coupled to HRP. Based on immunoelectrophoresis and/or ELISA, Anti-Mouse IgM(μ chain specific), AlpSdAbs® VHH(HRP) detects the μ chain of mouse IgM selectively, no reactivity with mouse, rabbit, human, cynomolgus, rat, goat IgG.

## Background

Most monoclonal antibodies are generated in mouse. There are five antibody isotypes (IgA, IgD, IgE, IgG, and IgM) from mouse. Each isotype has a different heavy chain. IgM accounts for 5-10% of the immunoglobulin pool and is the predominant antibody in the primary immune response. Unlike IgG, IgM does not contain a hinge region but does contain an additional constant domain. The monomeric form IgM has a molecular weight of 180 KD. It is classically represented as a pentamer of the basic four chain structure held together by a J chain but can also exist in a hexameric form without the J chain and as a monomer on the surface of B-cells.

VHH are single-domain antibodies derived from the variable regions of heavy chain of Camelidae immunoglobulin. The size of VHH is extremely small(<15KDa) compared to other forms of antibody fragment, which significantly increase the permeability of VHH. Thus VHH is considered of great value for research, diagnostics and therapeutics.

## Benefits


High lot-to-lot consistency  
Increased sensitivity and higher affinity  
Animal-free production

## Suggested Working Concentration


ELISA 1:5000-1:20000  
WB 1:5000-1:20000

Dilution factors are presented in the form of a range because the optimal dilution is a function of many factors, such as antigen density, permeability, etc. The actual dilution used must be determined empirically.

This product is for research use only and is not approved for use in humans or in clinical



A titer ELISA of mouse IgM. The plate was coated with different amounts of mouse IgM. 1:5000 dilution of Anti-Mouse IgM( $\mu$  chain specific), AlpSdAbs<sup>®</sup> VHH(HRP) was used as the detection antibody.



ELISA of specificity for different species of IgG. The plate was coated with 2ug/ml of different IgG. 1:1000 dilution of Anti-Mouse IgM( $\mu$  chain specific), AlpSdAbs<sup>®</sup> VHH(HRP) was used as the detection antibody.

This product is for research use only and is not approved for use in humans or in clinical