

# Anti-Chicken IgY, AlpSdAbs® VHH (HRP)

## Summary

|                           |   |
|---------------------------|---|
| Code                      | 024-101-005   |
| Immunogen                 | Recombinant Fc region of chicken IgY  |
| Host                      | Alpaca pacous   |
| Isotype                   | VHH domain of alpaca IgG2b/2c   |
| Conjugate                 | HRP   |
| Specificity               | Fc region of chicken IgY  |
| Cross-Reactivity          | No cross-reactivity with mouse, rabbit, human, cynomolgus, rat, goat IgG  |
| Purity                    | Recombinant Expression and Affinity purified  |
| Size                      | 100ug   |
| Buffer                    | 10mM PBS (pH 7.5), 0.05% sucrose, 0.1% trehalose, 0.01% proclin300  |
| Formation                 | Freeze-dried Powder, Store at 2-8 °C  |
| Rehydration               | Rehydrated by 100uL dH <sub>2</sub> O or 100uL dH <sub>2</sub> O with 50% Glycerol (Final concentration:1mg/mL) |
| Storage after Rehydration | -20 °C(Avoid freeze/thaw cycles), Stable for 12 months at -20°C, Protect from light                             |

## Description

Anti-Chicken IgY, AlpSdAbs® VHH(HRP) is designed for detecting chicken IgY specifically. Anti-Chicken IgY, AlpSdAbs® VHH(HRP) is based on monovalent, recombinant single domain antibodies to chicken IgY coupled to HRP. Based on immunoelectrophoresis and/or ELISA, Anti-Chicken IgY, AlpSdAbs® VHH(HRP) detects the chicken IgY selectively, no reactivity with mouse, rabbit, human, cynomolgus, rat, goat IgG.

## Background

Hens egg yolk immunoglobulins IgY can be transferred from the serum of mother hen to the offspring egg yolk to acquire immunity. In the immunodiagnostic technologies, IgY is an excellent antibody for using in immunological assays involving mammalian sera, due to discriminative properties of IgY compared to mammalian IgG, as IgY does not react with the rheumatoid factor and human anti-mouse IgG antibodies do not activate the complement system and do not bind to Fc receptor. Also, they have poor cross reactivity to mammalian IgG due to immunological differences. IgY does not contain a hinge region but does contain an additional constant domain. The whole IgY molecule possesses both the Fc region and the Fab region, which possessing the epitope-recognition site. The IgY contains two heavy and light chains. The common IgY is monomeric with a molecular weight of approximately 170 kDa.


VHH are single-domain antibodies derived from the variable regions of heavy chain of Camelidae immunoglobulin. The size of VHH is extremely small(<15kDa) compared to other forms of antibody fragment, which significantly increase the permeability of VHH. Thus VHH is considered of great value for research, diagnostics and therapeutics.

## Benefits


## Suggested Working Concentration

|   |  |                 |
|---|--|-----------------|
| High lot-to-lot consistency               | ELISA  | 1:10000-1:50000 |
| Increased sensitivity and higher affinity | WB   | 1:10000-1:50000 |
| Animal-free production                    | Dilution factors are presented in the form of a range because the optimal dilution is a function of many factors, such as antigen density, permeability, etc. The actual dilution used must be determined empirically. |                 |

This product is for research use only and is not approved for use in humans or in clinical



A titer ELISA of chicken IgY. The plate was coated with different amounts of chicken IgY. 1:10000 dilution of Anti-Chicken IgY, AlpSdAbs® VHH (HRP) was used as the detection antibody.



ELISA of specificity for different species of IgG. The plate was coated with 2ug/ml of different IgG. 1:1000 dilution of Anti-Chicken IgY, AlpSdAbs® VHH (HRP) was used as the detection antibody.

This product is for research use only and is not approved for use in humans or in clinical