

Anti-HA tag, AlpHcAbs[®] Rabbit antibody

Summary

Code	003-201-001
Immunogen	HA tag fused KLH
Host	Alpaca pacous
Isotype	VHH domain of alpaca IgG2b/2c fused to rabbit IgG Fc(mutation)
Conjugate	Unconjugated
Specificity	HA tag(YPYDVPDYA)
Cross-Reactivity	Highly selective for HA tag sequence
Purity	Recombinant Expression and Affinity purified
Concentration	1mg/ml
Formation	Liquid, 10mM PBS (pH 7.5), 0.05% sucrose, 0.1% trehalose, 0.01% proclin300,50% Glycerol
Storage	Store at -20 °C, (Avoid freeze / thaw cycles), Stable for 12 months at -20°C

Description

Anti-HA tag, AlpHcAbs[®] Rabbit antibody is designed for detecting HA tag fusion proteins specifically. Anti-HA tag, AlpHcAbs[®] Rabbit antibody is a monoclonal, recombinant, rabbit IgG Fc fused single domain antibody. Based on western blot and ELISA, Anti-HA tag, AlpHcAbs[®] Rabbit antibody detects the HA tag selectively, no reactivity with other proteins.

Background

The HA tag is widely used for detecting, manipulating or purifying proteins. This peptide can be expressed and detected with the protein of interest as an amino-terminal or carboxy-terminal fusion. Because of its small size, HA tag is unlikely to affect the tagged protein's biochemical properties. HA tag is useful for the labeling and detection of proteins using immunoblotting, immunoprecipitation, and immunostaining techniques.

Using antibody with Fc(mutation), the background from Fc receptors will be eliminated.

Benefits

- High lot-to-lot consistency
- Increased sensitivity and higher affinity
- Animal-free production

Suggested Working Concentration

ELISA	1:5000 -1:20000
WB	1:5000 -1:20000
ICC/IF	1:200-1:1000
IP	2ug/sample
Flow Cyt	1:200-1:1000

Dilution factors are presented in the form of a range because the optimal dilution is a function of many factors, such as antigen density, permeability, etc. The actual dilution used must be determined empirically.

This product is for research use only and is not approved for use in humans or in clinical