

Anti-Human IgG Fc, AlpSdAbs® VHH(VcMMAE ×4)

Summary

Code	023-101-101
Immunogen	Recombinant Fc region of human IgG
Host	Alpaca pacous
Isotype	VHH domain of alpaca IgG2b/2c
Conjugate	VcMMAE(2 moles VcMMAE per mole VHH)
Specificity	Human IgG(Fcγ fragment specific)
Cross-Reactivity	Recognizes human IgG Fcγ fragment specifically, and reacts with cynomolgus IgG. No Cross-reactivity to rabbit , mouse, rat, goat IgG
Purity	Recombinant Expression and Affinity purified
Concentration	0.5mg/ml
Formation	Liquid, 10mM PBS (pH 7.4)
Storage	Store at -20 °C(Avoid freeze / thaw cycles)

Description

Anti-Human IgG Fc, AlpSdAbs® VHH(VcMMAE ×4) is designed for studying on the internalization of antibodies. Anti-Human IgG Fc, AlpSdAbs® VHH(VcMMAE ×4) is based on monoclonal, recombinant single domain antibody to human IgG Fc coupled to VcMMAE. Based on immunoelectrophoresis and/or ELISA, Anti-Human IgG Fc, AlpSdAbs® VHH(VcMMAE ×4) reacts with the Fc fragment of human IgG heavy chain but not with the Fab portion of human IgG. Anti-Human IgG Fc, AlpSdAbs® VHH(VcMMAE ×4) is an effective detection tool and can be used as a useful tool for the evaluation of antibody potency prior to ADCs.

Background

VHH are single-domain antibodies derived from the variable regions of heavy chain of Camelidae immunoglobulin. The size of VHH is extremely small(<15KDa) compared to other forms of antibody fragment, which significantly increase the permeability of VHH. Thus VHH is considered of great value for research, diagnostics and therapeutics.

Benefits

High lot-to-lot consistency
 Increased sensitivity and higher affinity
 Animal-free production

Application notes

Antibody Internalization Test: 2ug per 10ug test antibody(molar ratio=2:1).

Dilution factors are presented in the form of a range because the optimal dilution is a function of many factors, such as antigen density, permeability, etc. The actual dilution used must be determined empirically.

This product is for research use only and is not approved for use in humans or in clinical

1) On Day 1 of the formal experiment, it was confirmed that the cells to be tested are in good condition. After thorough dilution and mixing, the cells are seeded at 1000-5000 cells/well/100uL in a 96 well cell culture plate. The test cells counts need to be adjusted by the user based on the growth rate and characteristics of the selected cells.

*Different cells growth rates vary greatly, and it is recommended to lay the board at a density of around 50% for subsequent endocytosis/internalization and killing detection.

Mix the test antibody(mAb) directly with VHH-MMAE in a molar ratio of 1:2 and let them mix thoroughly at room temperature (20-37 °C) for 20 minutes:

*The buffer of the test antibody in this step will not affect the binding of VHH-MMAE to form a complex.

Based on the user's mAb-VHH-MMAE complex concentration, prepare the test antibody into a 30ug/mL (200nM) solution using cell culture medium and mix evenly.

Dilute mAb-VHH-MMAE complex in a gradient using culture medium;

4) Add 100ul of diluted mAb-VHH-MMAE complex to a 96 well cell culture plate, put plate back to the incubator, and culture under suitable conditions for 48-96 hours.

*There are certain differences in the detection and cultivation time of different cells, test antibody and drug.

*Users can design controls group according to experimental needs and actual situations.

1) Observe cells growth and remove the culture medium from the cell culture plate before testing.

2) Use ATP content measurement method (such as CTG) to detect cells proliferation and vitality, in order to determine antibody internalization. Please refer to the instruction manual of the corresponding reagent kit for specific operating steps.

The calculation method for antibody internalization rate is:

$$\text{Internalization rate of test antibody} = [(Ac - As) / Ac] \times 100\%;$$

As: Experimental group absorbance (Test antibody+internalization detection reagent group):

Ac: Control group absorbance (Isotype Control antibody+internalization detection reagent group).